



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

breed. Many similar cases have come to my notice, especially among the warblers.—J. CLAIRE WOOD, *Detroit, Michigan*.

**The Breeding Habits of *Empidonax virescens* in Connecticut.**—On June 2, 1906, I was out collecting, in Stamford, Conn., with Mr. W. H. Hoyt. We were searching for a Hooded Warbler's nest in a dense laurel brake on the bank of a stream, when Mr. Hoyt found a nest of the Green-crested Flycatcher. The nest was compactly built, resembling a rather shallow nest of the Red-eyed Vireo, and was suspended from a fork at the end of a bush sapling about eight feet from the ground. The nest contained three fresh eggs. Both birds were shot, and upon dissection it was evident that the set was complete.

We proceeded about a mile up the river and there found an apparently well established colony of the birds. We first found a nest at the end of a limb of a large hemlock tree. The nest was about eight feet from the ground and was composed of shreds of inner hemlock bark. The cup was very shallow, and while the mass was packed into a fairly solid nest, a considerable number of shreds hung down in festoons from the nest for eight or ten inches. The three eggs were slightly incubated, and the female was so bold in their defence, darting at us and striking our hands when we touched the nest, that we were finally obliged to kill her with a stick before we could take the eggs.

Within 200 yards of this nest, we saw two other pairs of these birds, evidently breeding, although we were unable to find the nests. We found, however, six old nests, of which two were in hemlocks and four in bushes. The nests are so characteristic in situation and structure that I think there can be little doubt that these old nests were also those of *Empidonax virescens*. The nests are so frail that I do not believe they could hold together more than two years, which would seem to imply that three or four pairs have bred in this colony for several years.

On June 7, 1906, I returned to make another search for the nests. I clearly located three pairs of the birds and finally found one nest. It was composed of dead grass and was so roughly thrown together, with the strings and ends hanging down so loosely, that I should never have guessed it was a new nest had I not seen the bird fly to it. It also contained three eggs, and the bird was quite wild and shy. The nest was, as usual, in a fork at the end of a beech limb, about nine feet from the ground.

There are two unreported records by local collectors of *Empidonax virescens* breeding in Stamford (one by Mr. W. H. Hoyt, and one by Mr. G. Rowell), but they were made a number of years ago, and were merely rare and irregular finds. These records apparently show that the bird breeds regularly in Stamford; and is probably very much commoner with us than has been hitherto supposed.—LOUIS N. PORTER, *Stamford, Conn.*

***Empidonax griseus* Brewster vs. *Empidonax canescens* Salvin and Godman.**—In 'The Auk' for January, 1904, p. 80, I published a note showing that the two names given above apply to the same bird and that